

23136  
**PCT**

Serial No. 10/516,521  
WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07K 147/05</b>		A2	(11) International Publication Number: <b>WO 00/22001</b> (43) International Publication Date: <b>20 April 2000 (20.04.00)</b>
(21) International Application Number: <b>PCT/EP99/07671</b> (22) International Filing Date: <b>13 October 1999 (13.10.99)</b> (30) Priority Data: <b>98402540.3</b> <b>13 October 1998 (13.10.98)</b> <b>EP</b> (71) Applicant (for all designated States except US): <b>SANOFI-SYNTHELABO [FR/FR]; 174, avenue de France, F-75013 Paris (FR).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>RENARD, Stéphane [FR/FR]; 290, avenue Napoléon Bonaparte, F-92500 Rueil Malmaison (FR). AVENET, Patrick [FR/FR]; 50, rue de Vauhallan, F-91120 Palaiseau (FR).</b> (74) Agents: <b>ERNEST GUTMANN-YVES PLASSERAUD S.A. et al.; 3, rue Chauveau-Lagarde, F-75008 Paris (FR).</b>			(81) Designated States: <b>AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: <b>A NEW POTASSIUM CHANNEL OF THE ERG FAMILY</b>			
(57) Abstract  Herg4 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing herg4 polypeptides and polynucleotides in the design of protocols for the treatment of (i) dysfunctions of diseases, including, but not limited to, epilepsy, migraine, cell proliferation, comportmental troubles, among others and diagnostic assays for such conditions.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## A new potassium channel of the erg family

### FIELD OF INVENTION

This invention relates to newly identified  
5 polynucleotides, polypeptides encoded by such  
polynucleotides, to the use of such polynucleotides and  
polypeptides and to their production. More particularly, the  
polynucleotides and polypeptides of the present invention  
relate to erg potassium channel family, hereinafter referred  
10 to as herg4 (Human Erg Related Gene 4). The invention also  
relates to inhibiting or activating the action of such  
polynucleotides and polypeptides.

### BACKGROUND OF THE INVENTION

15 Potassium channels are involved in the maintenance  
of the cell resting membrane potential, and particularly for  
excitable cells in the control of their excitability by  
maintaining or allowing to recover the membrane potential.  
Potassium channels have also been involved in the regulation  
20 of cell proliferation.

This indicates that these channels have an  
interesting potential as therapeutic targets. Clearly there  
is a need for identification and characterization of further  
channels which can play a role in (i) preventing,  
25 ameliorating or correcting dysfunctions or diseases,  
including, but not limited to, epilepsy, migraine, cell  
proliferation, behavioural troubles and (ii) altering or  
preventing effect of endogenous neurotransmitters and  
hormones.

30

### SUMMARY OF THE INVENTION

In one aspect, the invention relates to herg4  
polypeptides and recombinant materials and methods for their  
production. Another aspect of the invention relates to  
35 methods for using such herg4 polypeptides and  
polynucleotides. Such uses include the treatment of  
epilepsy, migraine, cell proliferation, comportemental

troubles, to alter or prevent effect of endogenous neurotransmitters and hormones, or to diagnose or treat any disorder related to abnormal expression of these herg4 polypeptides, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with herg4 imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate herg4 activity or levels.

## DESCRIPTION OF THE INVENTION

### Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein-below.

"herg4" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2, or an allelic variant thereof.

"Receptor Activity" or "Channel Activity" or "Biological Activity of the Receptor" or "Biological Activity of the Channel" refers to the metabolic or physiologic function of said herg4 including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said herg4.

"herg4 gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:1 or allelic variants thereof and/or their complements.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of a Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not

"isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any  
5 polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA,  
10 and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions  
15 comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases  
20 such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and  
25 cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide  
30 isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include  
35 amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in

more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference

polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. See, e.g.: (COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or

polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., *Nucleic Acids Research* (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al., *J Molec Biol* (1990) 215:403).

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 or 3 terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical



to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

#### **Polypeptides of the Invention**

In one aspect, the present invention relates to herg4 polypeptides. The herg4 polypeptides include the polypeptide of SEQ ID NO:2, as well as polypeptides comprising the amino acid sequence of SEQ ID NO:2, and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Also included within herg4 polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO: 2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Preferably herg4 polypeptides exhibit at least one biological activity of the receptor.

The herg4 polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such

as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an  
5 additional sequence for stability during recombinant production.

Fragments of the herg4 polypeptides are also included in the invention. A fragment is a polypeptide having an amino acid sequence that entirely is the same as  
10 part, but not all, of the amino acid sequence of the aforementioned herg4 polypeptides. As with herg4 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region.  
15 Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of herg4 polypeptide. In this context "about" includes the particularly recited ranges larger or smaller  
20 by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of herg4 polypeptides, except for deletion of a continuous series of  
25 residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred are fragments characterized by structural or  
30 functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic  
35 regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate

receptor or channel activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain the biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

The herg4 polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

#### **Polynucleotides of the Invention**

Another aspect of the invention relates to herg4 polynucleotides. herg4 polynucleotides include isolated polynucleotides which encode the herg4 polypeptides and fragments, and polynucleotides closely related thereto. More specifically, herg4 polynucleotides of the invention include a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:1 encoding a herg4 polypeptide of SEQ ID NO: 2, and polynucleotide having the particular sequence of SEQ ID NO:1. herg4 polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity to a nucleotide sequence encoding the

herg4 polypeptide of SEQ ID NO:2 over its entire length, and a polynucleotide that is at least 80% identical to that having SEQ ID NO:1 over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under hERG4 polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:1 or contained in the cDNA insert in the plasmid deposited with the ATCC Deposit number 203078 to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, hERG4 polynucleotide includes nucleotide sequences having at least 80% identity to a nucleotide sequence encoding the hERG4 polypeptide expressed by the cDNA insert deposited at the ATCC with Deposit Number 203078, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above hERG4 polynucleotides.

A deposit containing a human hERG4 cDNA has been deposited with the American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA, on July 20, 1998, and assigned ATCC Deposit Number 203078. The deposited material (clone) is a DH5-a strain containing the expression vector pCR2.1 (Invitrogen) that further contains the hERG4 cDNA, referred to as "hERG4" upon deposit. The cDNA insert is within EcoRI site(s) in the vector. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the

event of any conflict with any description of sequences herein.

The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent.

herg4 of the invention is structurally related to other proteins of the potassium channel family, as shown by the results of sequencing the cDNA of SEQ ID NO:1 encoding human hERG4. The cDNA sequence of SEQ ID NO:1 contains an open reading frame (nucleotide number 1 to 3252) encoding a polypeptide of 1084 amino acids of SEQ ID NO:2. Amino acid sequence of SEQ ID NO:2 has about 35.2% identity (using BestFit version 1.0) in 885 amino acid residues with the drosophila elk protein (accession number: U04246). Nucleotide sequence of SEQ ID NO:1 has about 65% identity (using blastN version 1.4, GCG program package) in 2106 (197 to 2151) nucleotide residues with the elk cDNA (accession number: DMU04246).

One polynucleotide of the present invention encoding hERG4 may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in brain using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding hERG4 polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in SEQ ID NO:1 (nucleotide number 1 to 3252 of SEQ ID NO:1), or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

When the polynucleotides of the invention are used for the recombinant production of herg4 polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz et al., *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding herg4 variants comprising the amino acid sequence of herg4 polypeptide of SEQ ID NO:2 in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1 or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC with Deposit Number 203078 or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding herg4 and to

isolate cDNA and genomic clones of other genes that have a high sequence similarity to the herg4 gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding herg4 polypeptide comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO: 1 or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or alternatively conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease, either as a bulk composition or incorporated in a diagnostic kit.

30

### **Vectors, Host Cells, Expression**

The present invention also relates to a DNA or RNA molecule comprising an expression system, or vectors, wherein said expression system is capable of producing a herg4 polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO: 2 when said expression system is present in a compatible host cell. It also relates to host cells which

are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived  
5 from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells  
10 can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY* (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as  
15 calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include  
20 bacterial cells, such as streptococci, staphylococci, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and  
25 Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons,  
30 from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof,  
35 such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector



suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL* (supra).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the herg4 polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If herg4 polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

herg4 polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

### Diagnostic Assays

This invention also relates to diagnostic reagents comprising herg4 polynucleotides or fragments thereof, and their use in a diagnostic assay. It also relates to  
5 diagnostic kits containing such reagents for the detection of a mutated form of herg4 gene containing a polynucleotide of the invention. Detection of a mutated form of herg4 gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or  
10 susceptibility to a disease which results from under-expression, over-expression or altered expression of herg4. Individuals carrying mutations in the herg4 gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a  
15 subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion.  
20 Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled herg4 nucleotide sequences, such as described in Picketts et al, 1992, Human Genetics 89: 155-  
25 157.

In another embodiment, perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in  
30 electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., Science (1985) 230:1242. Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection  
35 or the chemical cleavage method. See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401. In another embodiment, an array of oligonucleotide probes comprising herg4 nucleotide sequence or fragments thereof can be

constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene  
5 expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996)).

The invention also relates to a diagnostic kit for the detection of a disease or a susceptibility to a disease  
10 in a subject related to expression or activity of herg4 polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO: 2 over its entire length in a subject containing a monoclonal or polyclonal antibody specific to herg4  
15 polypeptide, or a fragment thereof provided this fragment crossreact with monoclonal or polyclonal antibodies specific to herg4 polypeptide.

The diagnostic assays offer a process for diagnosing or determining a susceptibility to epilepsy, migraine, gliomas, abnormal proliferation and cancer, through  
20 detection of mutation in the herg4 gene by the methods described.

In addition, epilepsy, migraine, gliomas, abnormal proliferation and cancer, can be diagnosed by methods  
25 comprising determining from a sample derived from a subject an abnormally decreased or increased level of herg4 polypeptide or herg4 mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of  
30 polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a herg4, in a sample derived from a host are well-known to those of skill in the art.  
35 Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

### Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

### Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the herg4 polypeptides. The term "immunospecific" means that the antibodies have substantial greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the herg4 polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line

cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole et al., *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against herg4 polypeptides may also be employed to treat cerebral and cardiac and renal ischemias, brain and cardiac diseases, inflammation, pain, to mimic or antagonize effect of endogenous neurotransmitters and hormones, or to diagnose or treat any disorder related to abnormal expression of said herg4 polypeptides, among others.

25

#### **Vaccines/immunological products**

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with herg4 polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from epilepsy, migraine, to alter or prevent effect of endogenous neurotransmitters and hormones, or to diagnose or treat any disorder related to abnormal expression of the herg4 polypeptide, among others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering herg4 polypeptide via a vector directing expression of herg4

polynucleotide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a herg4 polypeptide wherein the composition comprises a herg4 polypeptide or herg4 gene. The vaccine formulation may further comprise a suitable carrier. Since herg4 polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

### Screening Assays

The herg4 polypeptide of the present invention may be employed in a screening process for compounds which bind the channel and which activate (agonists) or inhibit activation of (antagonists) the channel polypeptide of the present invention. The present invention also relates to screening assay for identifying agonists to herg4 polypeptide comprising an amino acid sequence which is at

least 80% identical to the amino acid sequence of  
SEQU ID NO: 2 over its entire length or a fragment thereof  
comprising:

(a) contacting host cells according to the invention  
5 with a candidate compound; and

(b) determining whether the candidate compound  
effects a signal generated by activation of the herg4  
polypeptide.

It also relates to screening assay for identifying  
10 antagonists to herg4 polypeptide comprising an amino acid  
sequence which is at least 80% identical to the amino acid  
sequence of SEQ ID NO: 2 over its entire length or a  
fragment thereof comprising:

(a) contacting said cell according to the invention  
15 with an agonist; and

(b) determining whether the signal generated by said  
agonist is diminished in the presence of a candidate  
compound.

Agonists and antagonists identified by such  
20 screening assays are encompassed in the scope of the  
invention.

Thus, polypeptides of the invention or their  
fragments may also be used to assess the binding of small  
molecule substrates and ligands in, for example, cells,  
25 cell-free preparations, chemical libraries, and natural  
product mixtures. These substrates and ligands may be  
natural substrates and ligands or may be structural or  
functional mimetics. See Coligan et al., *Current Protocols*  
*in Immunology* 1(2):Chapter 5 (1991).

30 herg4 polypeptides are implicated in many biological  
functions, and possibly pathologies. Accordingly, it is  
desirous to find compounds and drugs which stimulate herg4  
on the one hand and which can inhibit the function of herg4  
on the other hand. In general, agonists are employed for  
35 therapeutic and prophylactic purposes for such conditions as  
to alter effect of endogenous neurotransmitters and  
hormones. Antagonists may be employed for a variety of  
therapeutic and prophylactic purposes for such conditions as

epilepsy (hyperexcitability), migraines, abnormal proliferation and cancer, to antagonize effect of endogenous neurotransmitters and hormones.

5 In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a  
10 test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the channel is detected by means of a label directly or indirectly  
15 associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the channel, using detection systems appropriate to the cells bearing the  
20 channel at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Standard methods for conducting such screening assays are well understood in the  
25 art.

The recording of *herg4* channel activity may be carried out either by membrane voltage analysis of transfected cells or microinjected *xenopus* oocytes, directly (patch-clamp for example) or indirectly (fluorescent probes  
30 sensitive to changes of membrane potential). The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the channel is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with  
35 a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the channel, using detection systems appropriate to the cells bearing the channel at their



surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Standard methods for conducting such screening assays are well understood in the art.

Examples of potential herg4 antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the herg4, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the channel is prevented.

#### **Prophylactic and Therapeutic Methods**

This invention provides methods of treating abnormal conditions related to both an excess of and insufficient amounts of herg4 activity.

It also relates to therapeutic compositions for the treatment of a subject in need of enhanced activity or expression of herg4 polypeptide, containing:

(a) a therapeutically effective amount of an agonist of herg4 polypeptide according to the present invention, and/or

(b) a polynucleotide according to the invention in a form so as to effect production of said herg4 polypeptide activity in vivo, or a DNA or RNA vectors or a host cell according to the invention.

If the activity of herg4 is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as hereinabove described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking binding of ligands to the herg4, or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of herg4 polypeptides still capable of binding the ligand in competition with endogenous herg4 may be administered.

Typical embodiments of such competitors comprise fragments of the herg4 polypeptide.

In still another approach, expression of the gene encoding endogenous herg4 can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. See, for example, O'Connor, *J Neurochem* (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Alternatively, oligonucleotides which form triple helices with the gene can be supplied. See, for example, Lee et al., *Nucleic Acids Res* (1979) 6:3073; Cooney et al., *Science* (1988) 241:456; Dervan et al., *Science* (1991) 251:1360. These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

The invention also relates to a therapeutic composition for the treatment of a subject having need to inhibit activity or expression of herg4 polypeptide, containing:

- (a) a therapeutically effective amount of an antagonist of herg4 polypeptide according to the present invention, and/or
- (b) a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said herg4 polypeptide, and/or
- (c) a therapeutically effective amount of a polypeptide that competes with said herg4 polypeptide.

For treating abnormal conditions related to an under-expression of herg4 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates herg4, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of herg4 by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication

defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

#### 15 **Formulation and Administration**

Peptides, such as the soluble form of herg4 polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

30 Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants

such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or  
5 localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages,  
10 however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to  
15 require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated  
20 endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a  
25 retroviral plasmid vector. The cells are then introduced into the subject.

### Examples

The examples and figures below are carried out using  
30 standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples illustrate, but do not limit the invention.

Figure 1 is a photograph showing the results of a  
35 tissue distribution assay of herg4 protein using a Multiple Tissue Northern blot;

Figure 2 is a photograph showing the results of a tissue distribution assay of hERG4 protein using a Human Brain blot;

Figure 3 is an histogram showing the results of a tissue distribution assay of hERG4 protein using total mRNA samples from fifty human tissues. The numbers in abscissa has the following significance:

Tissues represented in figure 3:

1. Whole brain	16. Aorta	32. Liver	48. Fetal thymus
1. Amygdala	17. Skeletal muscle	33. Small Intestine	49. Fetal lung
2. Caudate nucleus	18. Colon	34. Spleen	50. Yeast tot RNA 0.1µg
3. Cerebellum	19. Bladder	35. Thymus	51. Yeast tRNA 0.1 µg
4. Cerebral cortex	20. Uterus	36. Peripheral leukocyte	52. E. Coli rRNA 0.1µg
5. Frontal lobe	21. Prostate	37. Lymph node	53. E. Coli DNA 0.1µg
6. Hippocampus	22. Stomach	38. Bone marrow	54. Poly r(A) 0.1µg
7. Medulla oblongata	23. Testis	39. Appendix	55. Human Cot-1 DNA 0.1µg
8. Occipital lobe	24. Ovary	40. Lung	56. Human DNA 0.1µg
9. Putamen	25. Pancreas	41. Trachea	57. Human DNA 0.5µg
10. Substantia nigra	26. Pituitary gland	42. Placenta	
11. Temporal lobe	27. Adrenal gland	43. Fetal brain	
12. Thalamus	28. Thyroid gland	44. Fetal heart	
13. Subthalamic nucleus	29. Salivary gland	45. Fetal Kidney	
14. Spinal cord	30. Mammary gland	46. Fetal liver	
15. Heart	31. Kidney	47. Fetal spleen	

Figure 4 is a graph showing the currents induced by CHO cells transfected with pcDNA3-hERG4 construct, in response to various depolarizations of the cells.

#### Example 1

#### Cloning the Human hERG4 cation Channel

The sequence of the hERG4 cation channel was first identified by searching a database containing approximately 2 million human ESTs, which was generated using high throughput automated DNA sequence analysis of randomly selected human cDNA clones (Adams, M.D. et al., Nature

377:3-174 (1995); Adams, M.D. et al., *Nature* 355:632-634 (1992); and Adams, M.D. et al., *Science* 252:1651-1656 (1991)). Sequence homology comparisons of each EST were performed against the GenBank database using the blastn and tblastn algorithms (Altschul, S.F. et al., *J. Mol. Biol.* 215:403-410 (1990)). A specific homology search using a cAMP binding domain of a cyclic nucleotide gated channel (CNG) amino acid sequence against this human EST database revealed one EST, from a cerebellum cDNA library, with approximately 60% similarity to CNG3. The sequence comparison suggested that it contained the partial open reading frame of a new protein. Sequence of the gene was confirmed by double strand DNA sequencing using the TaqFs (Perkin Elmer) and the gene was shown to be completely new by a blast search against Genbank release 110. The entire herg4 coding region was obtained with a combination of 5' sequence finding techniques and fragments were combined by recombinant DNA techniques, it was then amplified by PCR and inserted into the expression vector pCR2.1 (Invitrogen).

## Example 2

### Cloning and Expression of herg4 in Mammalian Cells

A typical mammalian expression vector contains the promoter element, which mediates the initiation of transcription of mRNA, the protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use in practicing the present invention include, for example, vectors such as PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109) and pcDNA3

(Invitrogen). Mammalian host cells that could be used include, human HEK 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and mouse L cells and Chinese hamster ovary (CHO) cells.

5 Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, zeocin or hygromycin allows the identification and isolation of the transfected cells.

10 The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection  
15 marker is the enzyme glutamine synthase (GS) (Murphy et al., *Biochem. J.* 227:277-279 (1991); Bebbington et al., *Bio/Technology* 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines  
20 contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

The expression vector pCMVSPORT3.0 contains the strong promoter (CMV) of the Cytomegalovirus. Multiple  
25 cloning sites, e.g., with the restriction enzyme cleavage site EcoRI, facilitate the cloning of the gene of interest.

### Example 3

#### Tissue distribution of herg4 mRNA expression

30 Northern blot analysis can be carried out to examine herg4 gene expression in human tissues, using methods described by, among others, Sambrook et al., cited above. A 1600 bp cDNA probe is obtained from the herg4 clone by EcorI-XhoI digestion and is radiolabelled with [<sup>32</sup>P]dATP.  
35 After labeling, the probe can be purified using a CHROMA SPIN- 100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified

labeled probe was then used to examine various human tissues for herg4 mRNA.

This probe is used to hybridize 3 membranes representing a population of messenger RNAs from different tissues in an Express Hybridization buffer (Clontech) according to the manufacturer specifications and protocol member PT1190-1, washed to a final stringency of 0.1 X SSC 0.1 %SDS 55°C. Following hybridization and washing, the blots can be mounted and the signal is detected using a Storm (Molecular Dynamics) apparatus after a 3 day exposure.

A first blot is carried out by using a Multiple Tissue Northern blot (MTN) (Clontech, ref. 7760-1), (2µg/lane), for assaying distribution of herg4 protein in human heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas mRNA. Results show that herg4 mRNA is distributed in brain (Figure 1).

A second blot is carried out by using a Human Brain blot (Clontech, ref. 7755-1) (2µg/lane) with cerebellum, cerebral cortex, medulla, spinal cord, occipital pole, frontal lobe, temporal lobe, putamen mRNA. Results show that herg4 is mainly distributed in cerebral cortex, temporal lobe, frontal lobe, occipital pole and putamen (Figure 2).

Another dot blot made with 100-500 ng total RNA samples from 50 human tissues (Clontech, ref. 7770-1), shows that herg4 messenger RNA is transcribed in the anterior brain (Figure 3). The localisation of herg4 pinpoints the role the protein of the invention can play in some physiological or pathophysiological mechanisms taking place in the anterior brain.

#### **Functional expression**

herg4 was expressed by transfection of the pcDNA3-herg4 construct in CHO cells. Currents were elicited by a depolarisation of the transfected CHO cells, obtained by applying a potential jump from -90mV to various potentials ranging from -70mV to + 95mv. Such a protocol is indicative of what happens when an action potential propagates along the neuron. Results show that Erg4 expresses a strong



current in response to the various potentials applied. The inversion potential of these currents is typical of a potassium channel. Queue currents are observed during repolarisation (figure 4).

5 Erg4 activates when the cell is depolarised and the resulting outward potassium current has the effect of repolarising the cell. As appears on figure 4, the current does not inactivate quickly. Thus, the cell is stably repolarised and subsequent action potentials would be  
10 inhibited. Such currents may obviously play an important role to reduce neuronal excitability.

#### Chromosomal localisation

A clone containing an exon of hERG4 is identified in a BAC genomic DNA library (genome systems, Inc) by a PCR  
15 with 2 sets of hERG4 specific primers. DNA from this clone is radiolabelled with digoxigenin dUTP by nick translation and the labelled probe is hybridized to normal metaphase chromosomes derived from PHA stimulated peripheral blood lymphocytes in a solution containing 50% formamide, 10%  
20 dextran and 2X SSC, and then cohybridized with a biotin labelled probe specific for the centromere of chromosome 11. This experiment has shown that the hERG4 gene is located in an area that corresponds to band 11p15..4 to 11p15..5 of chromosome 11. This locus is close to numerous genes  
25 important for brain function, in particular the locus for neuronal ceroid lipofuscinosis, a genetic form of epilepsy (Sharp et al, Human Molecular Genetics, 1997, vol. 6 n° 4 pp 591-5), and the locus for some human malignant astrocytomas (Fults et al, genomics 14 pp 799-801).

30 It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible in light of the above teachings and,  
35 therefore, are within the scope of the appended claims.

The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory

manuals, books, or other documents) cited herein are incorporated by reference.

## CLAIMS

1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity to a nucleotide sequence encoding the herg4 polypeptide of SEQ ID NO:2 over its entire length; or a nucleotide sequence complementary to said nucleotide sequence at least 80 % of said nucleotide sequence.

2. The polynucleotide of claim 1 which is DNA or RNA.

3. A polynucleotide according to one of claims 1 and 2 wherein said nucleotide sequence is at least 80% identical to that contained in SEQ ID NO:1.

4. A polynucleotide according to one of claims 1 and 3 wherein said nucleotide sequence comprises the herg4 polypeptide encoding sequence contained in SEQ ID NO:1.

5. A polynucleotide according to one of claims 1 and 4 which is polynucleotide of SEQ ID NO: 1.

6. An isolated herg4 polynucleotide comprising a nucleotide sequence selected from the group consisting of :

(a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the herg4 polypeptide expressed by the cDNA insert deposited at the ATCC with Deposit Number 203078 ; and

(b) a nucleotide sequence complementary to the nucleotide sequence of (a).

7. A nucleic acid probe for the detection of a polynucleotide according to claim 1 containing all or part of the complementary sequence of SEQ ID No. 1.

8. A DNA or RNA molecule comprising an expression system, or vectors, wherein said expression system is capable of producing a herg4 polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:2 when said expression system is present in a compatible host cell.

9.A host cell comprising the expression system of claim 8.

10. A herg4 polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:2 over its entire length.

11. The polypeptide of claim 11 which comprises the amino acid sequence of SEQ ID NO:2.

12. An antibody immunospecific for the herg4 polypeptide of claim 11.

13. Diagnostic kit for the detection of a mutated form of herg 4 gene containing a polynucleotide according to anyone of claims 1 to 7.

14. Therapeutic composition for the treatment of a subject in need of enhanced activity or expression of herg4 polypeptide, containing:

(a) a therapeutically effective amount of an agonist of herg4 polypeptide of claim 10, and/or

(b) a polynucleotide according to one of claims 1 to 6 in a form so as to effect production of said herg4 polypeptide activity in vivo, or a DNA or RNA according to claim 8 or a host cell according to claim 9.

15. Therapeutic composition for the treatment of a subject having need to inhibit activity or expression of herg4 polypeptide, containing:

(a) a therapeutically effective amount of an antagonist of herg4 polypeptide of claim 10, and/or

(b) a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said herg4 polypeptide, and/or

(c) a therapeutically effective amount of a polypeptide that competes with said herg4 polypeptide.

16. Diagnostic kit for the detection of a disease or a susceptibility to a disease in a subject related to expression or activity of herg4 polypeptide of claim 10 in a subject containing a monoclonal or polyclonal antibody specific to herg 4 polypeptide.

17. Screening assay for identifying agonists to herg4 polypeptide of claim 10 comprising:

(a) contacting host cells according to claim 9 with a candidate compound; and

5 (b) determining whether the candidate compound effects a signal generated by activation of the herg4 polypeptide.

18. An agonist identified by the assay of claim 17.

10 19. Screening assay for identifying antagonists to herg4 polypeptide of claim 10 comprising:

(a) contacting said cell according to claim 9 with an agonist; and

15 (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.

20. An antagonist identified by the assay of claim 19.

Figure 1 :




Figure 2 :




Figure 3





Figure 4





## SEQUENCE LISTING

5  
 <110> SANOFI-SYNTHELABO  
 <120> A new potassium channel of the erg family  
 10  
 <130> herg4  
 <140>  
 <141>  
 15  
 <160> 2  
 <170> PatentIn Ver. 2.1  
 20  
 <210> 1  
 <211> 3252  
 <212> DNA  
 <213> Homo sapiens  
 25  
 <220>  
 <221> CDS  
 <222> (1)..(3249)  
 <400> 1  
 30  
 atg ccg gcc atg cgg ggc ctc ctg gcg ccg cag aac acc ttc ctg gac 48  
 Met Pro Ala Met Arg Gly Leu Leu Ala Pro Gln Asn Thr Phe Leu Asp  
 1 5 10 15  
 acc atc gct acg cgc ttc gac ggc acg cac agt aac ttc gtg ctg ggc 96  
 35 Thr Ile Ala Thr Arg Phe Asp Gly Thr His Ser Asn Phe Val Leu Gly  
 20 25 30  
 aac gcc cag gtg gcg ggg ctc ttc ccc gtg gtc tac tgc tct gat ggc 144  
 40 Asn Ala Gln Val Ala Gly Leu Phe Pro Val Val Tyr Cys Ser Asp Gly  
 35 40 45  
 ttc tgt gac ctc acg ggc ttc tcc cgg gct gag gtc atg cag cgg ggc 192  
 Phe Cys Asp Leu Thr Gly Phe Ser Arg Ala Glu Val Met Gln Arg Gly  
 50 55 60  
 45  
 tgt gcc tgc tcc ttc ctt tat ggg cca gac acc agt gag ctc gtc cgc 240  
 Cys Ala Cys Ser Phe Leu Tyr Gly Pro Asp Thr Ser Glu Leu Val Arg  
 65 70 75 80  
 50  
 caa cag atc cgc aag gcc ctg gac gag cac aag gag ttc aag gct gag 288  
 Gln Gln Ile Arg Lys Ala Leu Asp Glu His Lys Glu Phe Lys Ala Glu  
 85 90 95  
 ctg atc ctg tac cgg aag agc ggg ctc ccg ttc tgg tgt ctc ctg gat 336  
 55 Leu Ile Leu Tyr Arg Lys Ser Gly Leu Pro Phe Trp Cys Leu Leu Asp  
 100 105 110  
 gtg ata ccc ata aag aat gag aaa ggg gag gtg gct ctc ttc cta gtc 384  
 60 Val Ile Pro Ile Lys Asn Glu Lys Gly Glu Val Ala Leu Phe Leu Val  
 115 120 125  
 tct cac aag gac atc agc gaa acc aag aac cga ggg ggc ccc gac aga 432  
 Ser His Lys Asp Ile Ser Glu Thr Lys Asn Arg Gly Gly Pro Asp Arg  
 130 135 140

	tgg	aag	gag	aca	ggt	ggt	ggc	cgg	cgc	cga	tat	ggc	cgg	gca	cga	tcc	480
	Trp	Lys	Glu	Thr	Gly	Gly	Gly	Arg	Arg	Arg	Tyr	Gly	Arg	Ala	Arg	Ser	
	145					150					155					160	
5	aaa	ggc	ttc	aat	gcc	aac	cgg	cgg	cgg	agg	cgg	gcc	gtg	ctc	tac	cac	528
	Lys	Gly	Phe	Asn	Ala	Asn	Arg	Arg	Arg	Ser	Arg	Ala	Val	Leu	Tyr	His	
					165					170					175		
10	ctg	tcc	ggg	cac	ctg	cag	aag	cag	ccc	aag	ggc	aag	cac	aag	ctc	aat	576
	Leu	Ser	Gly	His	Leu	Gln	Lys	Gln	Pro	Lys	Gly	Lys	His	Lys	Leu	Asn	
				180					185					190			
15	aag	ggg	gtg	ttt	ggg	gag	aaa	cca	aac	ttg	cct	gag	tac	aaa	gta	gcc	624
	Lys	Gly	Val	Phe	Gly	Glu	Lys	Pro	Asn	Leu	Pro	Glu	Tyr	Lys	Val	Ala	
			195					200					205				
20	gcc	atc	cgg	aag	tcg	ccc	ttc	atc	ctg	ttg	cac	tgt	ggg	gca	ctg	aga	672
	Ala	Ile	Arg	Lys	Ser	Pro	Phe	Ile	Leu	Leu	His	Cys	Gly	Ala	Leu	Arg	
		210					215					220					
25	gcc	acc	tgg	gat	ggc	ttc	atc	ctg	ctc	gcc	aca	ctc	tat	gtg	gct	gtc	720
	Ala	Thr	Trp	Asp	Gly	Phe	Ile	Leu	Leu	Ala	Thr	Leu	Tyr	Val	Ala	Val	
	225					230					235					240	
30	act	gtg	ccc	tac	agc	gtg	tgt	gtg	agc	aca	gca	cgg	gag	ccc	agt	gcc	768
	Thr	Val	Pro	Tyr	Ser	Val	Cys	Val	Ser	Thr	Ala	Arg	Glu	Pro	Ser	Ala	
					245					250					255		
35	gcc	cgc	ggc	ccg	ccc	agc	gtc	tgt	gac	ctg	gcc	gtg	gag	gtc	ctc	ttc	816
	Ala	Arg	Gly	Pro	Pro	Ser	Val	Cys	Asp	Leu	Ala	Val	Glu	Val	Leu	Phe	
				260					265					270			
40	atc	ctt	gac	att	gtg	ctg	aat	ttc	cgt	acc	aca	ttc	gtg	tcc	aag	tcg	864
	Ile	Leu	Asp	Ile	Val	Leu	Asn	Phe	Arg	Thr	Thr	Phe	Val	Ser	Lys	Ser	
			275					280					285				
45	ggc	cag	gtg	gtg	ttt	gcc	cca	aag	tcc	att	tgc	ctc	cac	tac	gtc	acc	912
	Gly	Gln	Val	Val	Phe	Ala	Pro	Lys	Ser	Ile	Cys	Leu	His	Tyr	Val	Thr	
		290					295					300					
50	acc	tgg	ttc	ctg	ctg	gat	gtc	atc	gcg	gcg	ctg	ccc	ttt	gac	ctg	cta	960
	Thr	Trp	Phe	Leu	Leu	Asp	Val	Ile	Ala	Ala	Leu	Pro	Phe	Asp	Leu	Leu	
	305					310					315					320	
55	cat	gcc	ttc	aag	gtc	aac	gtg	tac	ttc	ggg	gcc	cat	ctg	ctg	aag	acg	1008
	His	Ala	Phe	Lys	Val	Asn	Val	Tyr	Phe	Gly	Ala	His	Leu	Leu	Lys	Thr	
					325					330					335		
60	gtg	cgc	ctg	ctg	cgc	ctg	ctg	cgc	ctg	ctt	ccg	cgg	ctg	gac	cgg	tac	1056
	Val	Arg	Leu	Leu	Arg												

																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											</
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	----

	aag gcc aat gcc gac gtg aag ggg ctg acg tac tgc gtc ctg cag tgt	2016
	Lys Ala Asn Ala Asp Val Lys Gly Leu Thr Tyr Cys Val Leu Gln Cys	
	660 665 670	
5	ctg cag ctg gct ggc ctg cac gac agc ctt gcg ctg tac ccc gag ttt	2064
	Leu Gln Leu Ala Gly Leu His Asp Ser Leu Ala Leu Tyr Pro Glu Phe	
	675 680 685	
10	gcc ccg cgc ttc agt cgt ggc ctc cga ggg gag ctc agc tac aac ctg	2112
	Ala Pro Arg Phe Ser Arg Gly Leu Arg Gly Glu Leu Ser Tyr Asn Leu	
	690 695 700	
15	ggt gct ggg gga ggc tct gca gag gtg gac acc agc tcc ctg agc ggc	2160
	Gly Ala Gly Gly Gly Ser Ala Glu Val Asp Thr Ser Ser Leu Ser Gly	
	705 710 715 720	
20	gac aat acc ctt atg tcc acg ctg gag gag aag gag aca gat ggg gag	2208
	Asp Asn Thr Leu Met Ser Thr Leu Glu Glu Lys Glu Thr Asp Gly Glu	
	725 730 735	
25	cag ggc ccc acg gtc tcc cca gcc cca gct gat gag ccc tcc agc ccc	2256
	Gln Gly Pro Thr Val Ser Pro Ala Pro Ala Asp Glu Pro Ser Ser Pro	
	740 745 750	
30	ctg ctg tcc cct ggc tgc acc tcc tca tcc tca gct gcc aag ctg cta	2304
	Leu Leu Ser Pro Gly Cys Thr Ser Ser Ser Ser Ala Ala Lys Leu Leu	
	755 760 765	
35	tcc cca cgt cga aca gca ccc cgg cct cgt cta ggt ggc aga ggg agg	2352
	Ser Pro Arg Arg Thr Ala Pro Arg Pro Arg Leu Gly Gly Arg Gly Arg	
	770 775 780	
40	cca ggc agg gca ggg gct ttg aag gct gag gct ggc ccc tct gct ccc	2400
	Pro Gly Arg Ala Gly Ala Leu Lys Ala Glu Ala Gly Pro Ser Ala Pro	
	785 790 795 800	
45	cca cgg gcc cta gag ggg cta cgg ctg ccc ccc atg cca tgg aat gtg	2448
	Pro Arg Ala Leu Glu Gly Leu Arg Leu Pro Pro Met Pro Trp Asn Val	
	805 810 815	
50	ccc cca gat ctg agc ccc agg gta gta gat ggc att gaa gac ggc tgt	2496
	Pro Pro Asp Leu Ser Pro Arg Val Val Asp Gly Ile Glu Asp Gly Cys	
	820 825 830	
55	ggc tcg gac cag ccc aag ttc tct ttc cgc gtg ggc cag tct ggc ccg	2544
	Gly Ser Asp Gln Pro Lys Phe Ser Phe Arg Val Gly Gln Ser Gly Pro	
	835 840 845	
60	gaa tgt agc agc agc ccc tcc cct gga cca gag agc ggc ctg ctc act	2592
	Glu Cys Ser Ser Ser Pro Ser Pro Gly Pro Glu Ser Gly Leu Leu Thr	
	850 855 860	
65	gtt ccc cat ggg ccc agc gag gca agg aac aca gac aca ctg gac aag	2640
	Val Pro His Gly Pro Ser Glu Ala Arg Asn Thr Asp Thr Leu Asp Lys	
	865 870 875 880	
70	ctt cgg cag gcg gtg aca gag ctg tca gag cag gtg ctg cag atg cgg	2688
	Leu Arg Gln Ala Val Thr Glu Leu Ser Glu Gln Val Leu Gln Met Arg	
	885 890 895	
75	gaa gga ctg cag tca ctt cgc cag gct gtg cag ctt gtc ctg gcg ccc	2736
	Glu Gly Leu Gln Ser Leu Arg Gln Ala Val Gln Leu Val Leu Ala Pro	
	900 905 910	

	cac agg gag ggt ccg tgc cct cgg gca tgc gga gag ggg ccg tgc cca	2784
	His Arg Glu Gly Pro Cys Pro Arg Ala Ser Gly Glu Gly Pro Cys Pro	
	915 920 925	
5	gcc agc acc tcc ggg ctt ctg cag cct ctg tgt ctc gac act ggg gca	2832
	Ala Ser Thr Ser Gly Leu Leu Gln Pro Leu Cys Leu Asp Thr Gly Ala	
	930 935 940	
10	tcc tcc tac tgc ctg cag ccc cca gct ggc tct gtc ttg agt ggg act	2880
	Ser Ser Tyr Cys Leu Gln Pro Pro Ala Gly Ser Val Leu Ser Gly Thr	
	945 950 955 960	
15	tgg ccc cac cct cgt ccg ggg cct cct ccc ctc atg gca ccc tgg ccc	2928
	Trp Pro His Pro Arg Pro Gly Pro Pro Pro Leu Met Ala Pro Trp Pro	
	965 970 975	
20	tgg ggt ccc cca gcg tct cag agc tcc ccc tgg cct cga gcc aca gct	2976
	Trp Gly Pro Pro Ala Ser Gln Ser Ser Pro Trp Pro Arg Ala Thr Ala	
	980 985 990	
25	ttc tgg acc tcc acc tca gac tca gag ccc cct gcc tca gga gac ctc	3024
	Phe Trp Thr Ser Thr Ser Asp Ser Glu Pro Pro Ala Ser Gly Asp Leu	
	995 1000 1005	
30	tgc tct gag ccc agc acc cct gcc tcc cct cct cct tct gag gaa ggg	3072
	Cys Ser Glu Pro Ser Thr Pro Ala Ser Pro Pro Pro Ser Glu Glu Gly	
	1010 1015 1020	
35	gct agg act ggg ccc gca gag cct gtg agc cag gct gag gct acc agc	3120
	Ala Arg Thr Gly Pro Ala Glu Pro Val Ser Gln Ala Glu Ala Thr Ser	
	1025 1030 1035 1040	
40	act gga gag ccc cca cca ggg tca ggg ggc ctg gcc ttg ccc tgg gac	3168
	Thr Gly Glu Pro Pro Gly Ser Gly Gly Leu Ala Leu Pro Trp Asp	
	1045 1050 1055	
45	ccc cac agc ctg gag atg gtg ctt att ggc tgc cat ggc tct ggc aca	3216
	Pro His Ser Leu Glu Met Val Leu Ile Gly Cys His Gly Ser Gly Thr	
	1060 1065 1070	
50	gtc cag tgg acc cag gaa gaa ggc aca ggg gtc tga	3252
	Val Gln Trp Thr Gln Glu Glu Gly Thr Gly Val	
	1075 1080	
55	<210> 2	
	<211> 1083	
	<212> PRT	
	<213> Homo sapiens	
60	<400> 2	
	Met Pro Ala Met Arg Gly Leu Leu Ala Pro Gln Asn Thr Phe Leu Asp	
	1 5 10 15	
65	Thr Ile Ala Thr Arg Phe Asp Gly Thr His Ser Asn Phe Val Leu Gly	
	20 25 30	
70	Asn Ala Gln Val Ala Gly Leu Phe Pro Val Val Tyr Cys Ser Asp Gly	
	35 40 45	
75	Phe Cys Asp Leu Thr Gly Phe Ser Arg Ala Glu Val Met Gln Arg Gly	
	50 55 60	
80	Cys Ala Cys Ser Phe Leu Tyr Gly Pro Asp Thr Ser Glu Leu Val Arg	
	65 70 75 80	

6

Gln Gln Ile Arg Lys Ala Leu Asp Glu His Lys Glu Phe Lys Ala Glu  
85 90 95

5 Leu Ile Leu Tyr Arg Lys Ser Gly Leu Pro Phe Trp Cys Leu Leu Asp  
100 105 110

Val Ile Pro Ile Lys Asn Glu Lys Gly Glu Val Ala Leu Phe Leu Val  
115 120 125

10 Ser His Lys Asp Ile Ser Glu Thr Lys Asn Arg Gly Gly Pro Asp Arg  
130 135 140

Trp Lys Glu Thr Gly Gly Gly Arg Arg Arg Tyr Gly Arg Ala Arg Ser  
145 150 155 160

15 Lys Gly Phe Asn Ala Asn Arg Arg Arg Ser Arg Ala Val Leu Tyr His  
165 170 175

20 Leu Ser Gly His Leu Gln Lys Gln Pro Lys Gly Lys His Lys Leu Asn  
180 185 190

Lys Gly Val Phe Gly Glu Lys Pro Asn Leu Pro Glu Tyr Lys Val Ala  
195 200 205

25 Ala Ile Arg Lys Ser Pro Phe Ile Leu Leu His Cys Gly Ala Leu Arg  
210 215 220

Ala Thr Trp Asp Gly Phe Ile Leu Leu Ala Thr Leu Tyr Val Ala Val  
225 230 235 240

30 Thr Val Pro Tyr Ser Val Cys Val Ser Thr Ala Arg Glu Pro Ser Ala  
245 250 255

35 Ala Arg Gly Pro Pro Ser Val Cys Asp Leu Ala Val Glu Val Leu Phe  
260 265 270

Ile Leu Asp Ile Val Leu Asn Phe Arg Thr Thr Phe Val Ser Lys Ser  
275 280 285

40 Gly Gln Val Val Phe Ala Pro Lys Ser Ile Cys Leu His Tyr Val Thr  
290 295 300

Thr Trp Phe Leu Leu Asp Val Ile Ala Ala Leu Pro Phe Asp Leu Leu  
305 310 315 320

45 His Ala Phe Lys Val Asn Val Tyr Phe Gly Ala His Leu Leu Lys Thr  
325 330 335

50 Val Arg Leu Leu Arg Leu Leu Arg Leu Leu Pro Arg Leu Asp Arg Tyr  
340 345 350

Ser Gln Tyr Ser Ala Val Val Leu Thr Leu Leu Met Ala Val Phe Ala  
355 360 365

55 Leu Leu Ala His Trp Val Ala Cys Val Trp Phe Tyr Ile Gly Gln Arg  
370 375 380

Glu Ile Glu Ser Ser Glu Ser Glu Leu Pro Glu Ile Gly Trp Leu Gln  
385 390 395 400

60 Glu Leu Ala Arg Arg Leu Glu Thr Pro Tyr Tyr Leu Val Gly Arg Arg  
405 410 415

65 Pro Ala Gly Gly Asn Ser Ser Gly Gln Ser Asp Asn Cys Ser Ser Ser  
420 425 430

7

Ser Glu Ala Asn Gly Thr Gly Leu Glu Leu Leu Gly Gly Pro Ser Leu  
 435 440 445

5 Arg Ser Ala Tyr Ile Thr Ser Leu Tyr Phe Ala Leu Ser Ser Leu Thr  
 450 455 460

Ser Val Gly Phe Gly Asn Val Ser Ala Asn Thr Asp Thr Glu Lys Ile  
 465 470 475 480

10 Phe Ser Ile Cys Thr Met Leu Ile Gly Ala Leu Met His Ala Val Val  
 485 490 495

Phe Gly Asn Val Thr Ala Ile Ile Gln Arg Met Tyr Ala Arg Arg Phe  
 500 505 510

15 Leu Tyr His Ser Arg Thr Arg Asp Leu Arg Asp Tyr Ile Arg Ile His  
 515 520 525

20 Arg Ile Pro Lys Pro Leu Lys Gln Arg Met Leu Glu Tyr Phe Gln Ala  
 530 535 540

Thr Trp Ala Val Asn Asn Gly Ile Asp Thr Thr Glu Leu Leu Gln Ser  
 545 550 555 560

25 Leu Pro Asp Glu Leu Arg Ala Asp Ile Ala Met His Leu His Lys Glu  
 565 570 575

Val Leu Gln Leu Pro Leu Phe Glu Ala Ala Ser Arg Gly Cys Leu Arg  
 580 585 590

30 Ala Leu Ser Leu Ala Leu Arg Pro Ala Phe Cys Thr Pro Gly Glu Tyr  
 595 600 605

35 Leu Ile His Gln Gly Asp Ala Leu Gln Ala Leu Tyr Phe Val Cys Ser  
 610 615 620

Gly Phe Met Glu Val Leu Lys Gly Gly Thr Val Leu Ala Ile Leu Gly  
 625 630 635 640

40 Lys Gly Asp Leu Ile Gly Cys Glu Leu Pro Arg Arg Glu Gln Val Val  
 645 650 655

Lys Ala Asn Ala Asp Val Lys Gly Leu Thr Tyr Cys Val Leu Gln Cys  
 660 665 670

45 Leu Gln Leu Ala Gly Leu His Asp Ser Leu Ala Leu Tyr Pro Glu Phe  
 675 680 685

50 Ala Pro Arg Phe Ser Arg Gly Leu Arg Gly Glu Leu Ser Tyr Asn Leu  
 690 695 700

Gly Ala Gly Gly Gly Ser Ala Glu Val Asp Thr Ser Ser Leu Ser Gly  
 705 710 715 720

55 Asp Asn Thr Leu Met Ser Thr Leu Glu Glu Lys Glu Thr Asp Gly Glu  
 725 730 735

Gln Gly Pro Thr Val Ser Pro Ala Pro Ala Asp Glu Pro Ser Ser Pro  
 740 745 750

60 Leu Leu Ser Pro Gly Cys Thr Ser Ser Ser Ser Ala Ala Lys Leu Leu  
 755 760 765

65 Ser Pro Arg Arg Thr Ala Pro Arg Pro Arg Leu Gly Gly Arg Gly Arg  
 770 775 780

8

Pro Gly Arg Ala Gly Ala Leu Lys Ala Glu Ala Gly Pro Ser Ala Pro  
785 790 795 800

5 Pro Arg Ala Leu Glu Gly Leu Arg Leu Pro Pro Met Pro Trp Asn Val  
805 810 815

Pro Pro Asp Leu Ser Pro Arg Val Val Asp Gly Ile Glu Asp Gly Cys  
820 825 830

10 Gly Ser Asp Gln Pro Lys Phe Ser Phe Arg Val Gly Gln Ser Gly Pro  
835 840 845

15 Glu Cys Ser Ser Ser Pro Ser Pro Gly Pro Glu Ser Gly Leu Leu Thr  
850 855 860

Val Pro His Gly Pro Ser Glu Ala Arg Asn Thr Asp Thr Leu Asp Lys  
865 870 875 880

20 Leu Arg Gln Ala Val Thr Glu Leu Ser Glu Gln Val Leu Gln Met Arg  
885 890 895

Glu Gly Leu Gln Ser Leu Arg Gln Ala Val Gln Leu Val Leu Ala Pro  
900 905 910

25 His Arg Glu Gly Pro Cys Pro Arg Ala Ser Gly Glu Gly Pro Cys Pro  
915 920 925

30 Ala Ser Thr Ser Gly Leu Leu Gln Pro Leu Cys Leu Asp Thr Gly Ala  
930 935 940

Ser Ser Tyr Cys Leu Gln Pro Pro Ala Gly Ser Val Leu Ser Gly Thr  
945 950 955 960

35 Trp Pro His Pro Arg Pro Gly Pro Pro Pro Leu Met Ala Pro Trp Pro  
965 970 975

Trp Gly Pro Pro Ala Ser Gln Ser Ser Pro Trp Pro Arg Ala Thr Ala  
980 985 990

40 Phe Trp Thr Ser Thr Ser Asp Ser Glu Pro Pro Ala Ser Gly Asp Leu  
995 1000 1005

45 Cys Ser Glu Pro Ser Thr Pro Ala Ser Pro Pro Pro Ser Glu Glu Gly  
1010 1015 1020

Ala Arg Thr Gly Pro Ala Glu Pro Val Ser Gln Ala Glu Ala Thr Ser  
1025 1030 1035 1040

50 Thr Gly Glu Pro Pro Pro Gly Ser Gly Gly Leu Ala Leu Pro Trp Asp  
1045 1050 1055

Pro His Ser Leu Glu Met Val Leu Ile Gly Cys His Gly Ser Gly Thr  
1060 1065 1070

55 Val Gln Trp Thr Gln Glu Glu Gly Thr Gly Val  
1075 1080